

Continuous Birth Control Pill Use

Taking an active, hormone containing, pill every day is designed to stop all bleeding after an initial period of irregular bleeding. This handout explains how and gives tips to decrease the irregular bleeding.

Why do the spacer pills cause the uterus to bleed?

“The Period Pills,” “spacer,” “or “sugar” pills contain no active or hormone medication. The reason you bleed when you take spacer pills is because your hormone levels drop. You bleed because you did not take a progestin hormone or “real” birth control pill. The lining of the uterus needs stable hormone levels to prevent bleeding. The best way to prevent any bleeding or spotting is to have constant levels of the estrogen and the progesterone hormones, because these hormones support and keep the blood lining of the uterus stabilized.

What do birth control pills do to the uterus?

Birth control pills work to shrink the blood lining of the uterus. Over time the lining is so thin, the chances of unexpected bleeding and spotting become very low. It is very, very unlikely something is building up inside your uterus when you are on the pill. As a matter of fact, the risk of endometrial cancer decreases by 80% in women using the birth control pill for five years.

Irregular Bleeding is common at first

Break-through bleeding, or bleeding when you are not scheduled to bleed, is very common in the first 6 months of continuous birth control pill use. Your body is getting used to the constant level of hormones. Spotting is when the amount of blood is so tiny that no pad or tampon is needed. The longer you take the continuous pills the less bleeding and spotting will happen. You do not need to stop the pill to have a period because bleeding happens, instead try to figure out what caused the bleeding and keep taking the daily pill if you want to have no bleeding. Stopping the pill only begins the whole process again.

How can you help prevent a drop in the pill hormones and stop bleeding/spotting?

The most important thing is to take your pill as close as possible to the **same time every day**. Estrogen in the body begins to wear off, especially if you take your pill over 4 hours late.

Other suggestions if spotting continues:

All these suggestions and ideas listed below are to help you make it through the first six months of continuous pill use. Most women will have significantly less bleeding or spotting after six months. Keep a menstrual diary so you can learn what triggers a bleeding episode for you. Remember all women are individuals. You can learn about how you metabolize your pill and what works with your body.

- Alcohol:** Drinking alcohol keeps your liver busy detoxifying the alcohol so your hormone levels, especially estrogen, can be higher for a few days. If you drink everyday, even a glass of wine, your body could be used to the alcohol, so if you stop drinking, your estrogen levels may drop and trigger spotting.
- Tobacco:** Smoking can increase your metabolism of estrogen and result in lower levels of estrogen. If you smoke you now have another reason to quit or at least greatly reduce the amount you smoke.

- ❑ **Other medications:** Many medications, for example antibiotics, antifungals, anticonvulsants, and even herbal drugs like St. John's Wort, can change the amount of the pill hormones absorbed by the stomach and the metabolism of these hormones. It is very common to have some spotting with a new medication or a change in dose of medication. Sometimes these medications can actually decrease the pill hormones so much they become less effective at preventing pregnancy. Therefore, it is important to tell your provider about all the medications you are taking.
- ❑ **Time of day and stress** can affect your hormone levels. The progesterone receptors in the uterus look a little like cortisol receptors, so it might be possible that increased stress can trigger a change in progesterone activity. **Taking the pill at night**, before bed, could make the hormones peak when the cortisol levels are at nighttime levels and this could affect the activity of the hormones. Also, at night, the pill does not have to compete with food in your stomach to be absorbed. So, if you are having persistent spotting you could try switching the time of day you take your pill. However, you can expect some initial spotting with any change in the usual time you take your pill and it may take two weeks for your body to equilibrate to the new pill taking time.
- ❑ **Diarrhea or vomiting:** Anything that makes the pill go through your system too fast can make the pill not work as well because it was not absorbed or, worse, if it is lost in the vomit.
- ❑ **Altitude:** Some women report spotting when they take airplane trips or climb mountains. It could be the change in air pressure, just going to a new time zone, or even a change in your sleep patterns. If travelling in a different time zone, you should attempt to take your pill at the time based on your normal time zone.
- ❑ Non-steroidal anti-inflammatory medications, like **Naprosyn, Aspirin, or Ibuprofen** can decrease period bleeding and menstrual cramps, because they lessen the chemicals that cause period bleeding and decrease irritation in the lining of the uterus. Stop using them when your spotting stops. If your spotting continues after one week, you should call your provider, you may need a higher dose and your provider can give you a prescription. You should not use these drugs for more than 1-2 weeks or they could hurt your liver or kidneys.
- ❑ **Vitamin C**, 1000 mg, taken with your pill can help increase estrogen absorption for some women, so you should try this if the spotting has gone on for more than five days. However, you should stop taking the high dose of Vitamin C either when the spotting stops, or after a week if the spotting hasn't stopped. If you take it for too long, your body gets used to that large amount of Vitamin C, so that if you don't take it, you will then have a drop in estrogen levels and start spotting again!
- ❑ **Grapefruit juice** contains a chemical that slows estrogen metabolism if the pill is taken with a glass of juice. More estrogen may be available to your body to stop the spotting.

If you have any questions about any of these suggestions, please call your clinic. Often your provider can help and may even need to do an exam to find out why you are bleeding, because there may be an infection or change in health that is causing the bleeding. You can also email Dr. Miller at www.noperiod.com for suggestions. **Please call your clinic before you stop the birth control pill.**